



**KENTUCKY INJURY PREVENTION
AND RESEARCH CENTER**

Kentucky Resident Behavioral Health Indicators Report, 2016–2021

April 27, 2022

Prepared by

Lara Daniels (Lara.Daniels@uky.edu)
Ashley Bush
Mira Mirzaian

Released by

Kentucky Injury Prevention and Research Center (KIPRC)
333 Waller Avenue, Suite 242
Lexington, KY 40504
as bona fide agent for the Kentucky Department for Public Health

Suggested Citation

Daniels, L., Bush A., Mirzaian, M. (2022). *Kentucky Resident Behavioral Health Indicators Report, 2016–2021*.
Kentucky Injury Prevention and Research Center.

You may receive an email from KIPRC staff with a brief follow-up survey regarding this data report; thank you in advance for helping us to improve our services.



Kentucky Public Health
Prevent. Promote. Protect.

UK University of
Kentucky
College of Public Health

Table of Contents

Table 1. Numbers of Suicides among Youth and Adult Kentucky Residents: Demographics and Year, 2016–2021	3
Table 2. Numbers of Nonfatal Intentional Self-Harm Injury-Related Inpatient Hospitalizations among Youth and Adult Kentucky Residents: Demographics by Year, 2016–2021	4
Table 3. Numbers of Nonfatal Intentional Self-Harm Injury-Related Inpatient Hospitalizations among Youth and Adult Kentucky Residents by Diagnosis ¹ , Age Group, and Year 2016–2021.....	5
Table 4. Numbers of Nonfatal Intentional Self-Harm Injury-Related Emergency Department Visits among Youth and Adult Kentucky Residents: Demographics by Year, 2016–2021.....	6
Table 5. Numbers of Nonfatal Intentional Self-Harm Injury-Related Emergency Department Visits among Youth and Adult Kentucky Residents by Diagnosis ¹ , Age Group, and Year, 2016–2021	7
Table 6. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents: Demographics by Year, 2016–2021	8
Table 7. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents by Diagnosis ¹ , Age Group, and Year, 2016–2021.....	9
Table 8. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents: Demographics by Year, 2016–2021.....	10
Table 9. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents by Diagnosis ¹ , Age Group, and Year, 2016–2021	11
Table 10. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders associated with Substance Use among Kentucky Adult Residents: Demographics by Year, 2016–2021	12
Table 11. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders associated with Substance Use among Kentucky Adult Residents by Diagnosis ¹ and Year, 2016–2021.....	13
Table 12. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders associated with Substance Use among Kentucky Adult Residents: Demographics by Year, 2016–2021	14
Table 13. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders associated with Substance Use among Kentucky Adult Residents by Diagnosis ¹ and Year, 2016–2021.....	15
Table 14. Numbers of Nonfatal Drug Overdose Inpatient Hospitalizations among Kentucky Adult Residents: Demographics by Year, 2016–2021.....	16
Table 15. Numbers of Nonfatal Drug Overdose Inpatient Hospitalizations among Kentucky Adult Residents by Diagnosis ¹ and Year, 2016–2021	17
Table 16. Numbers of Nonfatal Drug Overdose Emergency Department Visits among Kentucky Adult Residents: Demographics by Year, 2016–2021.....	18
Table 17. Numbers of Nonfatal Drug Overdose Emergency Department Visits among Kentucky Adult Residents by Diagnosis ¹ and Year, 2016–2021.....	19
Definitions	20
Data Sources	20
Disclaimers	20

Table 1. Numbers of Suicides among Youth and Adult Kentucky Residents: Demographics and Year, 2016–2021

	Year						Total
	2016	2017	2018	2019	2020	2021	
Youth	25	24	21	23	22	23	138
Sex							
Female	*	*	6	8	*	9	42
Male	*	*	15	15	*	14	96
Adult	715	712	736	704	733	742	4,342
Race							
Black	32	28	32	34	40	36	202
White	669	679	690	658	682	690	4,068
Other	14	5	14	12	11	16	72
Sex							
Female	157	133	130	146	109	110	785
Male	558	579	606	558	624	632	3,557
Age Group							
18–24	73	65	80	66	82	87	453
25–34	119	118	112	120	129	131	729
35–44	125	130	123	141	142	134	795
45–54	153	146	147	116	115	123	800
55–64	115	116	117	136	123	101	708
65–74	61	91	91	83	71	97	494
75+	69	46	66	42	71	69	363
Total	740	736	757	727	755	765	4,480

Note:

*Due to state data release policy, counts in certain cells of the tables have been suppressed either because 1) the observed number of events is very small (a count of at least one but fewer than five) and not appropriate for publication or 2) it could be used to calculate the number in a cell that has been suppressed.

Youth are defined as Kentucky residents 17 years of age and younger. Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 2. Numbers of Nonfatal Intentional Self-Harm Injury-Related Inpatient Hospitalizations among Youth and Adult Kentucky Residents: Demographics by Year, 2016–2021

	Year						Total
	2016	2017	2018	2019	2020	2021	
Youth	303	342	390	423	435	559	2,452
Race							
Black	48	55	46	73	64	80	366
White	247	282	335	336	362	467	2,029
Other	8	5	9	14	9	12	57
Sex							
Female	248	291	320	321	357	454	1,991
Male	55	51	70	102	78	105	461
Age Group							
<14	72	58	89	99	122	149	589
14–17	231	284	301	324	313	410	1,863
Adult	2,853	2,867	2,704	2,590	2,376	2,161	15,551
Race							
Black	190	208	200	206	193	204	1,201
White	2,632	2,619	2,465	2,347	2,138	1,917	14,118
Other	31	40	39	37	45	40	232
Sex							
Female	1,645	1,659	1,515	1,395	1,312	1,181	8,707
Male	1,208	1,208	1,189	1,195	1,064	980	6,844
Age Group							
18–24	543	567	584	553	563	524	3,334
25–34	635	649	614	576	540	458	3,472
35–44	641	610	550	547	483	483	3,314
45–54	581	561	500	476	386	324	2,828
55–64	312	338	304	284	254	235	1,727
65–74	97	97	113	103	108	95	613
75+	44	45	39	51	42	42	263
Total	3,156	3,209	3,094	3,013	2,811	2,720	18,003

Note:

Youth are defined as Kentucky residents 17 years of age and younger. Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 3. Numbers of Nonfatal Intentional Self-Harm Injury-Related Inpatient Hospitalizations among Youth and Adult Kentucky Residents by Diagnosis¹, Age Group, and Year 2016–2021

Diagnosis	Year						Total
	2016	2017	2018	2019	2020	2021	
Youth							
Other	175	186	182	215	207	290	1,255
Poisoning by, adverse effect of, and underdosing of 4-aminophenol derivatives	32	46	51	58	63	91	341
Poisoning by other and unspecified antidepressants	30	39	47	44	70	68	298
Intentional self-harm by unspecified sharp object	23	11	41	39	43	48	205
Poisoning by, adverse effect of, and underdosing of other nonsteroidal anti-inflammatory drugs [NSAIDs]	24	36	32	31	27	43	193
Poisoning by, adverse effect of, and underdosing of other and unspecified drugs, medicaments, and biological substances	19	24	37	36	25	19	160
Adults							
Other	1,928	1,893	1,778	1,685	1,521	1,403	10,208
Poisoning by, adverse effect of, and underdosing of other and unspecified antipsychotics and neuroleptics	187	218	208	200	186	142	1,141
Poisoning by, adverse effect of, and underdosing of benzodiazepines	244	247	186	167	149	152	1,145
Poisoning by other and unspecified antidepressants	201	212	211	179	174	162	1,139
Intentional self-harm by unspecified sharp object	147	136	166	194	169	168	980
Poisoning by, adverse effect of, and underdosing of other and unspecified drugs, medicaments, and biological substances	146	161	155	165	177	134	938

Note:

¹Diagnosis based on first listed intentional self-harm injury ICD-10-CM code.

Youth are defined as Kentucky residents 17 years of age and younger. Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 4. Numbers of Nonfatal Intentional Self-Harm Injury-Related Emergency Department Visits among Youth and Adult Kentucky Residents: Demographics by Year, 2016–2021

	Year						Total
	2016	2017	2018	2019	2020	2021	
Youth	986	1,132	1,056	1,017	924	1,203	6,318
Race							
Black	82	103	91	85	104	132	597
White	883	999	934	905	795	1,020	5,536
Other	21	30	31	27	25	51	185
Sex							
Female	745	859	803	767	719	926	4,819
Male	241	273	253	250	205	277	1,499
Age Group							
<10	21	24	12	31	20	37	145
10–13	201	209	235	261	239	318	1,463
14–17	764	899	809	725	665	848	4,710
Adult	2,746	2,694	2,787	2,623	2,603	2,421	15,874
Race							
Black	244	265	276	270	316	296	1,667
White	2,476	2,392	2,472	2,315	2,243	2,070	13,968
Other	26	37	39	38	44	55	239
Sex							
Female	1,450	1,411	1,493	1,417	1,400	1,374	8,545
Male	1,296	1,283	1,294	1,206	1,203	1,047	7,329
Age Group							
18–24	813	790	904	804	854	819	4,984
25–34	739	806	785	741	706	631	4,408
35–44	621	531	538	519	503	490	3,202
45–54	376	368	348	371	336	268	1,925
55–64	146	153	139	137	144	134	995
65–74	38	30	58	37	45	57	265
75+	13	16	15	14	15	22	95
Total	3,732	3,826	3,843	3,640	3,527	3,624	22,192

Note:

Youth are defined as Kentucky residents 17 years of age and younger. Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 5. Numbers of Nonfatal Intentional Self-Harm Injury-Related Emergency Department Visits among Youth and Adult Kentucky Residents by Diagnosis¹, Age Group, and Year, 2016–2021

Diagnosis	Year						Total
	2016	2017	2018	2019	2020	2021	
Youth							
Other	455	502	518	497	481	672	3,125
Intentional self-harm by unspecified sharp object	144	172	162	134	112	139	863
Intentional self-harm by other sharp object	139	154	159	132	103	117	804
Intentional self-harm by other specified means	98	125	86	110	62	78	559
Poisoning by other antidepressants, intentional self-harm, initial encounter	85	103	70	77	89	117	541
Poisoning by, adverse effect of, and underdosing of other and unspecified drugs, medicaments, and biological substances	65	76	61	67	77	80	426
Adults							
Other	1,602	1,503	1,496	1,464	1,395	1,350	8,810
Intentional self-harm by unspecified sharp object	284	302	347	311	318	293	1,855
Intentional self-harm by other sharp object	235	254	255	254	239	204	1,441
Intentional self-harm by other specified means	215	234	278	232	246	192	1,397
Intentional self-harm by knife	238	231	257	211	234	186	1,357
Poisoning by, adverse effect of, and underdosing of other and unspecified drugs, medicaments, and biological substances	172	170	154	151	171	196	1,014

Note:

¹Diagnosis based on first listed intentional self-harm injury ICD-10-CM code.

Youth are defined as Kentucky residents 17 years of age and younger. Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 6. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents: Demographics by Year, 2016–2021

	Year						Total
	2016	2017	2018	2019	2020	2021	
Youth	1,642	1,703	1,740	1,893	1,548	1,795	10,321
Race							
Black	248	220	231	259	208	224	1,390
White	1,371	1,452	1,475	1,592	1,311	1,527	8,728
Other	23	31	34	42	29	44	203
Sex							
Female	1,088	1,161	1,145	1,284	1,074	1,273	7,025
Male	554	542	595	609	474	522	3,296
Age Group							
<10	96	49	19	14	20	20	218
10–13	430	419	487	595	441	551	2,923
14–17	1,116	1,235	1,234	1,284	1,087	1,224	7,180
Adult							
	17,696	17,741	15,542	14,492	13,244	12,950	91,665
Race							
Black	1,214	1,247	1,100	1,168	1,140	1,135	7,004
White	16,270	16,262	14,263	13,177	11,946	11,638	83,556
Other	212	232	179	147	158	177	1,105
Sex							
Female	9,397	9,451	8,193	7,590	6,769	6,660	48,060
Male	8,299	8,290	7,349	6,902	6,475	6,290	43,605
Age Group							
18–24	2,382	2,576	2,332	2,288	2,138	2,094	13,810
25–34	3,976	3,967	3,439	3,143	2,974	2,942	20,441
35–44	3,738	3,717	3,175	2,983	2,785	2,836	19,234
45–54	3,616	3,377	2,933	2,639	2,191	2,167	16,923
55–64	2,428	2,418	2,139	1,957	1,772	1,658	12,372
65–74	1,030	1,113	986	920	894	794	5,737
75+	526	573	538	562	490	459	3,148
Total	19,338	19,444	17,282	16,385	14,792	14,745	101,986

Note:

Youth are defined as Kentucky residents 17 years of age and younger. Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 7. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents by Diagnosis¹, Age Group, and Year, 2016–2021

Diagnosis	Year						Total
	2016	2017	2018	2019	2020	2021	
Youth							
Other	595	604	598	577	478	3,492	6,984
Major depressive disorder, recurrent severe without psychotic features	407	493	636	648	611	3,470	6,940
Major depressive disorder, single episode, unspecified	196	182	168	292	138	1,143	2,286
Unspecified mood [affective] disorder	301	179	109	138	102	929	1,858
Major depressive disorder, single episode, severe without psychotic features	68	94	93	104	104	541	1,082
Other persistent mood [affective] disorders	50	115	103	94	78	505	1,010
Adults							
Other	10,075	10,023	9,113	8,354	7,730	8,039	53,334
Major depressive disorder, recurrent severe without psychotic features	2,928	3,071	2,530	2,365	2,243	2,233	15,370
Major depressive disorder, single episode, unspecified	2,299	1,996	1,493	1,343	921	719	8,771
Bipolar disorder, unspecified	1,045	1,110	981	852	820	676	5,484
Major depressive disorder, recurrent, severe with psychotic symptoms	778	853	787	726	681	651	4,476
Adjustment disorders	533	635	604	822	830	601	4,025

Note:

¹Diagnosis based on first listed mental and behavioral disorder ICD-10-CM code.

Youth are defined as Kentucky residents 17 years of age and younger. Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change April 2022.

Table 8. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents: Demographics by Year, 2016–2021

	Year						Total
	2016	2017	2018	2019	2020	2021	
Youth	19,455	21,168	20,733	20,442	13,074	12,495	107,367
Race							
Black	2,986	3,361	3,302	3,254	1,725	1,495	16,123
White	16,117	17,416	17,030	16,773	11,069	10,718	89,123
Other	352	391	401	415	280	282	2,121
Sex							
Female	9,226	10,092	10,056	10,149	6,757	6,817	53,097
Male	10,229	11,076	10,677	10,293	6,317	5,678	54,270
Age Group							
<10	4,030	4,233	3,980	3,568	1,999	1,718	19,528
10–13	5,528	6,141	6,274	6,370	3,953	3,792	32,058
14–17	9,897	10,794	10,479	10,504	7,122	6,985	55,781
Adult	210,483	217,747	225,504	234,858	187,750	170,242	1,246,584
Race							
Black	19,404	22,361	23,828	26,069	19,873	16,841	128,376
White	189,497	193,616	199,614	206,441	165,879	151,311	1,106,358
Other	1,582	1,770	2,062	2,348	1,998	2,090	11,850
Sex							
Female	140,186	145,488	149,959	157,018	121,760	110,363	824,774
Male	70,297	72,259	75,545	77,840	65,990	59,879	421,810
Age Group							
18–24	22,647	23,588	23,995	25,561	20,287	18,649	134,727
25–34	40,931	41,623	42,299	42,863	34,511	30,637	232,864
35–44	42,560	43,640	44,223	45,383	36,860	32,814	245,480
45–54	41,887	42,463	43,415	43,363	34,766	30,408	236,302
55–64	30,900	32,747	34,497	36,707	29,213	26,670	190,734
65–74	17,598	18,962	20,794	22,799	18,486	17,971	116,610
75+	13,960	14,724	16,281	18,182	13,627	13,093	89,867
Total	229,938	238,915	246,237	255,300	200,824	182,737	1,353,951

Note:

Youth are defined as Kentucky residents 17 years of age and younger. Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 9. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents by Diagnosis¹, Age Group, and Year, 2016–2021

Diagnosis	Year						Total
	2016	2017	2018	2019	2020	2021	
Youth							
Attention-deficit/hyperactivity disorder, unspecified type	8,420	9,196	8,966	8,289	4,787	3,686	43,344
Other	4,703	4,954	4,595	4,545	3,233	4,170	26,200
Anxiety disorder, unspecified	2,494	3,001	3,059	3,293	2,132	2,256	15,967
Major depressive disorder, single episode, unspecified	2,605	2,697	2,934	3,162	2,313	1,753	15,732
Bipolar disorder, unspecified	698	716	670	563	303	286	3,236
Oppositional defiant disorder	535	604	509	590	306	344	2,888
Adults							
Anxiety disorder, unspecified	72,458	71,988	74,577	77,502	63,889	58,209	418,623
Major depressive disorder, single episode, unspecified	68,378	72,301	74,612	76,411	56,297	42,351	390,350
Other	36,501	40,038	41,556	43,886	36,730	42,677	241,388
Bipolar disorder, unspecified	17,586	18,125	17,498	18,330	14,256	11,720	97,515
Other specified anxiety disorders	7,441	8,487	8,262	9,828	9,552	9,342	52,912
Generalized anxiety disorder	8,119	6,808	8,999	8,901	7,026	5,943	45,796

Note:

¹Diagnosis based on first listed mental and behavioral disorder ICD-10-CM code.

Youth are defined as Kentucky residents 17 years of age and younger. Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 10. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders associated with Substance Use among Kentucky Adult Residents: Demographics by Year, 2016–2021

	Year						Total
	2016	2017	2018	2019	2020	2021	
Adult							
Race							
Black	361	401	401	357	429	574	2,523
White	6,587	7,396	6,589	5,563	5,405	5,846	37,386
Other	66	74	51	36	74	102	403
Female	2,490	2,729	2,424	1,904	1,736	1,980	13,263
Male	4,524	5,142	4,617	4,052	4,172	4,542	27,049
Age Group							
18–24	504	565	444	280	273	293	2,359
25–34	1,794	2,152	1,814	1,445	1,231	1,408	9,844
35–44	1,838	2,102	1,922	1,602	1,675	1,827	10,966
45–54	1,595	1,671	1,529	1,295	1,322	1,423	8,835
55–64	962	1,012	982	997	1,018	1,136	6,107
65–74	260	309	274	265	318	369	1,795
75+	61	60	76	72	71	66	406
Total	7,014	7,871	7,041	5,956	5,908	6,522	40,312

Note:

Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 11. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders associated with Substance Use among Kentucky Adult Residents by Diagnosis¹ and Year, 2016–2021

Diagnosis	Year						Total
	2016	2017	2018	2019	2020	2021	
Adult							
Alcohol dependence	3,079	3,365	3,193	3,211	3,460	3,613	19,921
Other	1,459	1,790	1,406	1,013	891	998	7,557
Opioid addiction and dependence	1,781	1,794	1,436	847	648	745	7,251
Alcohol abuse	313	322	327	259	388	623	2,232
Other stimulant abuse	170	286	392	363	335	337	1,883
Other psychoactive substance use, unspecified	212	314	287	263	186	206	1,468

Notes:

¹Diagnosis based on first listed mental and behavioral disorder due to substance use ICD-10-CM code.

Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 12. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders associated with Substance Use among Kentucky Adult Residents: Demographics by Year, 2016–2021

	Year						Total
	2016	2017	2018	2019	2020	2021	
Adult							
Race							
Black	8,458	8,711	8,755	10,066	9,386	9,681	55,057
White	51,677	53,051	56,107	57,759	54,711	54,521	327,826
Other	564	660	741	735	811	1,130	4,641
Sex							
Female	21,976	22,918	24,765	25,959	23,471	24,037	143,126
Male	38,723	39,504	40,838	42,601	41,437	41,295	244,398
Age Group							
18–24	7,462	7,336	7,135	6,977	6,571	6,693	42,174
25–34	15,948	16,713	17,464	17,844	16,737	16,133	100,839
35–44	13,982	14,727	16,069	17,602	17,161	17,475	97,016
45–54	12,770	12,418	12,630	12,813	12,067	11,894	74,592
55–64	7,559	8,269	8,944	9,729	8,943	9,098	52,542
65–74	2,321	2,321	2,592	2,740	2,622	3,228	15,824
75+	657	638	769	855	807	811	4,537
Total	60,699	62,422	65,603	68,560	64,908	65,332	387,524

Note:

Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 13. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders associated with Substance Use among Kentucky Adult Residents by Diagnosis¹ and Year, 2016–2021

Diagnosis	Year						Total
	2016	2017	2018	2019	2020	2021	
Adult							
Other	20,184	22,086	24,135	25,644	23,912	26,356	142,317
Alcohol abuse	17,492	16,489	16,367	17,127	15,199	13,647	96,321
Other psychoactive substance abuse	8,785	10,102	11,637	10,840	11,451	10,248	63,063
Alcohol dependence	6,513	5,901	6,657	6,797	6,177	6,017	38,062
Cannabis abuse	4,778	4,139	3,622	4,336	3,634	3,766	24,275
Alcohol abuse with intoxication, unspecified	2,947	3,705	3,185	3,816	4,535	5,298	23,486

Notes:

¹Diagnosis based on first listed mental and behavioral disorder due to substance use ICD-10-CM code.

Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 14. Numbers of Nonfatal Drug Overdose Inpatient Hospitalizations among Kentucky Adult Residents: Demographics by Year, 2016–2021

	Year						Total
	2016	2017	2018	2019	2020	2021	
Adult							
Race							
Black	410	427	412	360	380	427	2,416
White	5,382	5,222	4,511	4,100	4,058	3,624	26,897
Other	53	60	58	54	58	64	347
Sex							
Female	3,283	3,083	2,738	2,427	2,367	2,163	16,061
Male	2,562	2,626	2,243	2,087	2,129	1,952	13,599
Age Group							
18–24	631	594	548	551	559	552	3,405
25–34	1,046	1,097	915	851	873	738	5,520
35–44	1,129	1,113	979	846	918	851	5,836
45–54	1,370	1,261	1,051	948	887	715	5,678
55–64	834	830	711	639	646	672	4,886
65–74	503	470	459	381	375	374	2,562
75+	332	344	318	298	238	243	1,773
Total	5,845	5,709	4,981	4,514	4,496	4,115	29,660

Note:

Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 15. Numbers of Nonfatal Drug Overdose Inpatient Hospitalizations among Kentucky Adult Residents by Diagnosis¹ and Year, 2016–2021

Diagnosis	Year						Total
	2020	2019	2018	2017	2016	2021	
Adult							
Other	3,416	3,249	2,982	2,675	2,503	2,177	17,002
Poisoning by, adverse effect of, and underdosing of other and unspecified drugs, medicaments, and biological substances	689	678	533	498	593	586	3,577
Poisoning by, adverse effect of, and underdosing of benzodiazepines	662	620	495	409	385	375	2,946
Poisoning by, adverse effect of, and underdosing of psychostimulants	298	369	371	375	352	388	2,153
Poisoning by, adverse effect of, and underdosing of other opioids	396	362	330	290	334	300	2,012
Poisoning by, adverse effect of, and underdosing of narcotics and psychodysleptics	384	431	270	267	329	289	1,970

Notes:

¹Diagnosis based on first listed drug overdose ICD-10-CM code.

Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 16. Numbers of Nonfatal Drug Overdose Emergency Department Visits among Kentucky Adult Residents: Demographics by Year, 2016–2021

	Year						Total
	2016	2017	2018	2019	2020	2021	
Adult							
Race							
Black	785	857	885	839	1,150	1,115	5,631
White	9,732	10,670	8,573	8,170	9,380	9,226	55,751
Other	84	138	91	115	136	165	729
Sex							
Female	4,827	5,013	4,293	3,984	4,320	4,381	26,818
Male	5,774	6,652	5,256	5,140	6,346	6,125	35,293
Age Group							
18–24	1,723	1,674	1,405	1,221	1,436	1,425	8,884
25–34	3,580	4,008	3,190	2,998	3,374	3,031	20,181
35–44	2,360	2,703	2,167	2,208	2,752	2,793	14,983
45–54	1,545	1,819	1,449	1,369	1,610	1,542	8,704
55–64	698	764	684	695	866	1,018	5,355
65–74	395	394	364	336	380	435	2,304
75+	300	303	290	297	248	262	1,700
Total	10,601	11,665	9,549	9,124	10,666	10,506	62,111

Note:

Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2022.

Table 17. Numbers of Nonfatal Drug Overdose Emergency Department Visits among Kentucky Adult Residents by Diagnosis¹ and Year, 2016–2021

Diagnosis	Year						Total
	2016	2017	2018	2019	2020	2021	
Adult							
Poisoning by, adverse effect of, and underdosing of narcotics and psychodysleptics	4,388	5,077	3,485	3,136	3,571	2,611	22,268
Other	3,349	3,520	3,469	3,275	3,311	3,264	20,188
Poisoning by, adverse effect of, and underdosing of other and unspecified drugs, medicaments, and biological substances	1,183	1,314	1,115	1,166	1,501	1,851	8,130
Poisoning by, adverse effect of, and underdosing of other and unspecified narcotics	597	703	556	654	1,037	1,363	4,910
Poisoning by, adverse effect of, and underdosing of other opioids	472	517	458	514	803	970	3,734
Poisoning by, adverse effect of, and underdosing of benzodiazepines	612	534	466	379	443	447	2,881

Notes:

¹Diagnosis based on first listed drug overdose ICD-10-CM code.

Produced by the Kentucky Injury Prevention and Research Center, as bona fide agent for the Kentucky Department for Public Health. Data are provisional and subject to change. April 2021.

Definitions

Indicator	ICD-10/ICD-10-CM Codes	
	Emergency Department Visits	Inpatient Hospitalizations
Mental and behavioral disorders	F20–F69 or F90–F99 in any field	F20–F69 and F90–F99 in the primary diagnosis field
Mental and behavioral disorders due to substance use	F10–F16 or F18–F19 in any field	F10–F16 and F18–F19 in the primary diagnosis field
	Fatal	Nonfatal
Self-harm (from CSTE Injury Surveillance Toolkit)	Underlying cause of death code of X60–X84, Y87.0, or U03	In any field: X71–X83; T36–T50 with 6th character = 2 (Note: include T36.9, T37.9, T39.9, T41.4, T42.7, T43.9, T45.9, T47.9, and T49.9 with 5th character = 2); T51–T65 with 6th character = 2 (Note: include T51.9, T52.9, T53.9, T54.9, T56.9, T57.9, T58.0, T58.1, T58.9, T59.9, T60.9, T61.0, T61.1, T61.9, T62.9, T63.9, T64.0, T64.8, and T65.9 with a 5th character = 2); T71 with 6th character = 2; and T14.91; all codes must have a 7th character of A or missing, excluding adverse effects and underdosing
Drug overdose (from CSTE Injury Surveillance Toolkit)	Underlying cause of death code of X40–X44, X60–X64, X85, or Y10–Y14	In any field: T36–T50; for T36.9, T37.9, T39.9, T41.4, T42.7, T43.9, T45.9, T47.9, and T49.9, a 5th character; for all others, a 6th character; all codes must have a 7th character of A or missing, excluding adverse effects and underdosing

Data Sources

Emergency department visits: Kentucky Outpatient Services Database, Office of Health Data and Analytics, Cabinet for Health and Family Services

Inpatient hospitalizations: Kentucky Inpatient Hospitalization Claims Files, Office of Health Data and Analytics, Cabinet for Health and Family Services

Deaths: Kentucky Death Certificate Database, Kentucky Office of Vital Statistics, Cabinet for Health and Family Services

Disclaimers

*Due to state data release policy, counts in certain cells of the tables have been suppressed either because 1) the observed number of events is very small (a count of at least one but fewer than five) and not appropriate for publication or 2) it could be used to calculate the number in a cell that has been suppressed.

Emergency department visits and inpatient hospitalization counts represent encounters of care and could be greater than the total number of patients treated. To avoid duplicate counts, inpatient hospitalizations resulting in

death and emergency department visits resulting in inpatient hospitalization or death have been excluded from this analysis. Data are provisional and subject to change. Produced by the Kentucky Injury Prevention and Research Center as bona fide agent for the Kentucky Department for Public Health, April 2022. This report was supported by grants numbers NU17CE010064 (Centers for Disease Control and Prevention) and 3048114413. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention or the U.S. Department of Health and Human Services.