

Kentucky Resident Behavioral Health Indicators Report, 2017–2022

August 2023

Prepared by

Alaina Murphy (alaina.murphy@uky.edu) Lara Daniels

Released by

Kentucky Injury Prevention and Research Center (KIPRC) University of Kentucky 2536 Harrodsburg Road, Suite B340 Lexington, KY 40504 as bona fide agent for the Kentucky Department for Public Health

Suggested Citation

Murphy, A., Daniels, L., (2023). *Kentucky Resident Behavioral Health Indicators Report*, 2017–2022. Kentucky Injury Prevention and Research Center.

You may receive an email from KIPRC staff with a brief follow-up survey regarding this data report; thank you in advance for helping us to improve our services.





Table of Contents

| Table 1. Numbers of Suicides among Youth and Adult Kentucky Residents: Demographics and Year, 2017–2022 | 3 |
|--|----|
| Table 2. Numbers of Nonfatal Intentional Self-Harm Injury-Related Inpatient Hospitalizations amongYouth and Adult Kentucky Residents: Demographics by Year, 2017–2022 | 4 |
| Table 3. Numbers of Nonfatal Intentional Self-Harm Injury-Related Inpatient Hospitalizations amongYouth and Adult Kentucky Residents by Diagnosis ¹ , Age Group, and Year 2017–2022 | 5 |
| Table 4. Numbers of Nonfatal Intentional Self-Harm Injury-Related Emergency Department Visitsamong Youth and Adult Kentucky Residents: Demographics by Year, 2017–2022 | 6 |
| Table 5. Numbers of Nonfatal Intentional Self-Harm Injury-Related Emergency Department Visits among Youth and Adult Kentucky Residents by Diagnosis ¹ , Age Group, and Year, 2017–2022 | 7 |
| Table 6. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents: Demographics by Year, 2017–2022 | 8 |
| Table 7. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents by Diagnosis ¹ , Age Group, and Year, 2017–2022 | 9 |
| Table 8. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents: Demographics by Year, 2017–2022 | 10 |
| Table 9. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents by Diagnosis ¹ , Age Group, and Year, 2017–2022 | 11 |
| Table 10. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders Associated with Substance Use among Kentucky Adult Residents: Demographics by Year, 2017–2022 | 12 |
| Table 11. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders Associated with Substance Use among Kentucky Adult Residents by Diagnosis ¹ and Year, 2017–2022 | 13 |
| Table 12. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders Associated with Substance Use among Kentucky Adult Residents: Demographics by Year, 2017–2022 | |
| Table 13. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders Associated with Substance Use among Kentucky Adult Residents by Diagnosis ¹ and Year, 2017–2022 | |
| Table 14. Numbers of Nonfatal Drug Overdose Inpatient Hospitalizations among Kentucky Adult Residents: Demographics by Year, 2017–2022 | 16 |
| Table 15. Numbers of Nonfatal Drug Overdose Inpatient Hospitalizations among Kentucky Adult Residents by Diagnosis ¹ and Year, 2017–2022 | 17 |
| Table 16. Numbers of Nonfatal Drug Overdose Emergency Department Visits among Kentucky Adult Residents: Demographics by Year, 2017–2022 | 18 |
| Table 17. Numbers of Nonfatal Drug Overdose Emergency Department Visits among Kentucky Adult Residents by Diagnosis ¹ and Year, 2017–2022 | 19 |
| Definitions | 20 |
| Data Sources | 20 |
| Disclaimers | 20 |

| | | | Ye | ar | | | Total |
|-----------|------|------|------|------|------|------|-------|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | |
| Youth | 24 | 21 | 23 | 22 | 23 | 28 | 141 |
| Sex | | | | | | | |
| Female | 10 | 6 | 8 | 5 | 9 | 6 | 44 |
| Male | 14 | 15 | 15 | 17 | 14 | 22 | 97 |
| Adult | 712 | 736 | 704 | 733 | 740 | 761 | 4,386 |
| Race | | | | | | | |
| Black | 28 | 32 | 34 | 40 | 36 | 50 | 220 |
| White | 679 | 690 | 658 | 682 | 688 | 689 | 4,086 |
| Other | 5 | 14 | 12 | 11 | 16 | 22 | 80 |
| Sex | | | | | | | |
| Female | 133 | 130 | 146 | 109 | 108 | 136 | 762 |
| Male | 579 | 606 | 558 | 624 | 632 | 625 | 3,624 |
| Age Group | | | | | | | |
| 18–24 | 65 | 80 | 66 | 82 | 86 | 81 | 460 |
| 25–34 | 118 | 112 | 120 | 129 | 131 | 129 | 739 |
| 35–44 | 130 | 123 | 141 | 142 | 134 | 149 | 819 |
| 45–54 | 146 | 147 | 116 | 115 | 123 | 128 | 775 |
| 55–64 | 116 | 117 | 136 | 123 | 101 | 109 | 702 |
| 65–74 | 91 | 91 | 83 | 71 | 97 | 90 | 523 |
| 75+ | 46 | 66 | 42 | 71 | 68 | 75 | 368 |
| Total | 712 | 736 | 704 | 733 | 740 | 761 | 4,386 |

Table 1. Numbers of Suicides among Youth and Adult Kentucky Residents: Demographics and Year, 2017–2022

Note:

| | | | Ye | ear | | | Total |
|-----------|-------|-------|-------|-------|-------|-------|--------|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | |
| Youth | 342 | 390 | 423 | 435 | 559 | 494 | 2,643 |
| Race | | | | | | | |
| Black | 55 | 46 | 73 | 64 | 80 | 79 | 397 |
| White | 282 | 335 | 336 | 362 | 467 | 406 | 2,188 |
| Other | 5 | 9 | 14 | 9 | 12 | 9 | 58 |
| Sex | | | | | | | |
| Female | 291 | 320 | 321 | 357 | 454 | 404 | 2,147 |
| Male | 51 | 70 | 102 | 78 | 105 | 90 | 496 |
| Age Group | | | | | | | |
| <14 | 58 | 89 | 99 | 122 | 149 | 121 | 638 |
| 14–17 | 284 | 301 | 324 | 313 | 410 | 373 | 2,005 |
| Adult | 2,867 | 2,704 | 2,590 | 2,376 | 2,161 | 2,039 | 14,737 |
| Race | | | | | | | |
| Black | 208 | 200 | 206 | 193 | 204 | 185 | 1,196 |
| White | 2,619 | 2,465 | 2,347 | 2,138 | 1,917 | 1,804 | 13,290 |
| Other | 40 | 39 | 37 | 45 | 40 | 50 | 251 |
| Sex | | | | | | | |
| Female | 1,659 | 1,515 | 1,395 | 1,312 | 1,181 | 1,142 | 8,204 |
| Male | 1,208 | 1,189 | 1,195 | 1,064 | 980 | 897 | 6,533 |
| Age Group | | | | | | | |
| 18–24 | 567 | 584 | 553 | 563 | 524 | 425 | 3,216 |
| 25–34 | 649 | 614 | 576 | 540 | 458 | 461 | 3,298 |
| 35–44 | 610 | 550 | 547 | 483 | 483 | 425 | 3,098 |
| 45–54 | 561 | 500 | 476 | 386 | 324 | 341 | 2,588 |
| 55–64 | 338 | 304 | 284 | 254 | 235 | 234 | 1,649 |
| 65–74 | 97 | 113 | 103 | 108 | 95 | 112 | 628 |
| 75+ | 45 | 39 | 51 | 42 | 42 | 41 | 260 |
| Total | 3,209 | 3,094 | 3,013 | 2,811 | 2,720 | 2,533 | 17,380 |

Table 2. Numbers of Nonfatal Intentional Self-Harm Injury-Related Inpatient Hospitalizations among Youth and Adult Kentucky Residents: Demographics by Year, 2017–2022

Note:

Table 3. Numbers of Nonfatal Intentional Self-Harm Injury-Related Inpatient Hospitalizations among Youth and Adult Kentucky Residents by Diagnosis¹, Age Group, and Year 2017–2022

| Diagnosis | | | Yea | ar | | | Total |
|---|-------|-------|-------|-------|-------|-------|-------|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | |
| Youth | | | | | | | |
| Other | 186 | 182 | 215 | 207 | 290 | 243 | 1,323 |
| Poisoning by, adverse effect of, and underdosing of 4-aminophenol derivatives | 46 | 51 | 58 | 63 | 91 | 80 | 389 |
| Poisoning by other and unspecified antidepressants | 39 | 47 | 44 | 70 | 68 | 78 | 346 |
| Intentional self-harm by unspecified sharp object | 11 | 41 | 39 | 43 | 48 | 35 | 217 |
| Poisoning by, adverse effect of, and underdosing of other nonsteroidal anti-inflammatory drugs [NSAIDs] | 36 | 32 | 31 | 27 | 43 | 32 | 201 |
| Poisoning by, adverse effect of, and underdosing of other and unspecified drugs, medicaments, and biological substances | 24 | 37 | 36 | 25 | 19 | 26 | 167 |
| Adults | | | | | | | |
| Other | 1,893 | 1,778 | 1,685 | 1,521 | 1,403 | 1,292 | 9,572 |
| Poisoning by other and unspecified antidepressants | 212 | 211 | 179 | 174 | 162 | 186 | 1,124 |
| Poisoning by, adverse effect of, and underdosing of other and unspecified antipsychotics and neuroleptics | 218 | 208 | 200 | 186 | 142 | 154 | 1,108 |
| Poisoning by, adverse effect of, and underdosing of benzodiazepines | 247 | 186 | 167 | 149 | 152 | 127 | 1,028 |
| Intentional self-harm by unspecified sharp object | 136 | 166 | 194 | 169 | 168 | 122 | 955 |
| Poisoning by, adverse effect of, and underdosing of other and unspecified drugs, medicaments, and biological substances | 161 | 155 | 165 | 177 | 134 | 158 | 950 |

Note:

¹Diagnosis based on first listed intentional self-harm injury ICD-10-CM code.

| | | | Ye | ar | | | Total |
|-----------|-------|-------|-------|-------|-------|-------|--------|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | |
| Youth | 1,132 | 1,056 | 1,017 | 924 | 1,203 | 1,102 | 6,434 |
| Race | | | | | | | |
| Black | 103 | 91 | 85 | 104 | 132 | 106 | 621 |
| White | 999 | 934 | 905 | 795 | 1,020 | 960 | 5,613 |
| Other | 30 | 31 | 27 | 25 | 51 | 36 | 200 |
| Sex | | | | | | | |
| Female | 859 | 803 | 767 | 719 | 926 | 830 | 4,904 |
| Male | 273 | 253 | 250 | 205 | 277 | 272 | 1,530 |
| Age Group | | | | | | | |
| <10 | 24 | 12 | 31 | 20 | 37 | 26 | 150 |
| 10–13 | 209 | 235 | 261 | 239 | 318 | 305 | 1,567 |
| 14–17 | 899 | 809 | 725 | 665 | 848 | 771 | 4,717 |
| Adult | 2,694 | 2,787 | 2,623 | 2,603 | 2,421 | 2,219 | 15,347 |
| Race | | | | | | | |
| Black | 265 | 276 | 270 | 316 | 296 | 260 | 1,683 |
| White | 2,392 | 2,472 | 2,315 | 2,243 | 2,070 | 1,899 | 13,391 |
| Other | 37 | 39 | 38 | 44 | 55 | 60 | 273 |
| Sex | | | | | | | |
| Female | 1,411 | 1,493 | 1,417 | 1,400 | 1,374 | 1,230 | 8,325 |
| Male | 1,283 | 1,294 | 1,206 | 1,203 | 1,047 | 989 | 7,022 |
| Age Group | | | | | | | |
| 18–24 | 790 | 904 | 804 | 854 | 819 | 697 | 4,868 |
| 25–34 | 806 | 785 | 741 | 706 | 631 | 625 | 4,294 |
| 35–44 | 531 | 538 | 519 | 503 | 490 | 419 | 3,000 |
| 45–54 | 346 | 321 | 342 | 303 | 268 | 255 | 1,835 |
| 55–64 | 175 | 166 | 166 | 177 | 134 | 152 | 970 |
| 65–74 | 30 | 58 | 37 | 45 | 57 | 39 | 266 |
| 75+ | 16 | 15 | 14 | 15 | 22 | 32 | 114 |
| Total | 3,826 | 3,843 | 3,640 | 3,527 | 3,624 | 3,321 | 21,781 |

Table 4. Numbers of Nonfatal Intentional Self-Harm Injury-Related Emergency Department Visits among Youth and Adult Kentucky Residents: Demographics by Year, 2017–2022

Note:

| Diagnosis | | | Ye | ar | | | Total |
|---|-------|-------|-------|-------|-------|-------|-------|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | |
| Youth | | | | | | | |
| Other | 502 | 518 | 497 | 481 | 672 | 601 | 3,271 |
| Intentional self-harm by unspecified sharp object | 172 | 162 | 134 | 112 | 139 | 123 | 842 |
| Intentional self-harm by other sharp object | 154 | 159 | 132 | 103 | 117 | 102 | 767 |
| Poisoning by other antidepressants, intentional self-harm, initial encounter | 103 | 70 | 77 | 89 | 117 | 108 | 564 |
| Intentional self-harm by other specified means | 125 | 86 | 110 | 62 | 78 | 71 | 532 |
| Poisoning by, adverse effect of, and underdosing of other and unspecified drugs, medicaments, and biological substances | 76 | 61 | 67 | 77 | 80 | 97 | 458 |
| Adults | | | | | | | |
| Other | 1,503 | 1,496 | 1,464 | 1,395 | 1,350 | 1,256 | 8,464 |
| Intentional self-harm by unspecified sharp object | 302 | 347 | 311 | 318 | 293 | 252 | 1,823 |
| Intentional self-harm by other sharp object | 254 | 255 | 254 | 239 | 204 | 194 | 1,400 |
| Intentional self-harm by other specified means | 234 | 278 | 232 | 246 | 192 | 146 | 1,328 |
| Intentional self-harm by knife | 231 | 257 | 211 | 234 | 186 | 196 | 1,315 |
| Poisoning by, adverse effect of, and underdosing of other and unspecified drugs, medicaments, and biological substances | 170 | 154 | 151 | 171 | 196 | 175 | 1,017 |

Table 5. Numbers of Nonfatal Intentional Self-Harm Injury-Related Emergency Department Visits among Youth and Adult Kentucky Residents by Diagnosis¹, Age Group, and Year, 2017–2022

Note:

¹Diagnosis based on first listed intentional self-harm injury ICD-10-CM code.

| | | | Ye | or | | | |
|-----------|--------|--------|--------|--------|--------|--------|--------|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | Total |
| Youth | 1,703 | 1,740 | 1,893 | 1,548 | 1,795 | 1,817 | 10,496 |
| Race | 1,705 | 1,740 | 1,075 | 1,540 | 1,775 | 1,017 | 10,490 |
| Black | 220 | 231 | 259 | 208 | 224 | 222 | 1,364 |
| White | 1,452 | 1,475 | 1,592 | 1,311 | 1,527 | 1,565 | 8,922 |
| Other | 31 | 34 | 42 | 29 | 44 | 30 | 210 |
| Sex | | | | | | | |
| Female | 1,161 | 1,145 | 1,284 | 1,074 | 1,273 | 1,274 | 7,211 |
| Male | 542 | 595 | 609 | 474 | 522 | 543 | 3,285 |
| Age Group | | | | | | | |
| <10 | 49 | 19 | 14 | 20 | 20 | 20 | 142 |
| 10–13 | 419 | 487 | 595 | 441 | 551 | 540 | 3,033 |
| 14–17 | 1,235 | 1,234 | 1,284 | 1,087 | 1,224 | 1,257 | 7,321 |
| Adult | 17,741 | 15,542 | 14,492 | 13,244 | 12,950 | 11,607 | 85,576 |
| Race | | | | | | | |
| Black | 1,247 | 1,100 | 1,168 | 1,140 | 1,135 | 960 | 6,750 |
| White | 16,262 | 14,263 | 13,177 | 11,946 | 11,638 | 10,445 | 77,731 |
| Other | 232 | 179 | 147 | 158 | 177 | 202 | 1,095 |
| Sex | | | | | | | |
| Female | 9,451 | 8,193 | 7,590 | 6,769 | 6,660 | 5.959 | 44,622 |
| Male | 8,290 | 7,349 | 6,902 | 6,475 | 6,290 | 5,648 | 40,954 |
| Age Group | | | | | | | |
| 18–24 | 2,576 | 2,332 | 2,288 | 2,138 | 2,094 | 1,883 | 13,311 |
| 25–34 | 3,967 | 3,439 | 3,143 | 2,974 | 2,942 | 2,524 | 18,989 |
| 35–44 | 3,717 | 3,175 | 2,983 | 2,785 | 2,836 | 2,450 | 17,946 |
| 45–54 | 3,377 | 2,933 | 2,639 | 2,191 | 2,167 | 1,924 | 15,231 |
| 55–64 | 2,418 | 2,139 | 1,957 | 1,772 | 1,658 | 1,600 | 11,544 |
| 65–74 | 1,113 | 986 | 920 | 894 | 794 | 814 | 5,521 |
| 75+ | 573 | 538 | 562 | 490 | 459 | 412 | 3,034 |
| Total | 19,444 | 17,282 | 16,385 | 14,792 | 14,745 | 13,424 | 96,072 |

Table 6. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents: Demographics by Year, 2017–2022

Note:

| Diagnosis | | | Ye | ar | | | Total |
|---|--------|-------|-------|-------|-------|-------|--------|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | |
| Youth | | | | | | | |
| Other | 604 | 598 | 577 | 478 | 640 | 791 | 3,688 |
| Major depressive disorder, recurrent severe without psychotic features | 493 | 636 | 648 | 611 | 675 | 622 | 3,685 |
| Major depressive disorder, single episode, unspecified | 182 | 168 | 292 | 138 | 167 | 85 | 1,032 |
| Unspecified mood [affective] disorder | 179 | 109 | 138 | 102 | 100 | 100 | 728 |
| Major depressive disorder, single episode, severe without psychotic features | 94 | 93 | 104 | 104 | 78 | 81 | 554 |
| Other persistent mood [affective] disorders | 115 | 103 | 94 | 78 | 65 | 82 | 537 |
| Adults | | | | | | | |
| Other | 10,023 | 9,113 | 8,354 | 7,730 | 8,039 | 7,470 | 50,729 |
| Major depressive disorder, recurrent severe without psychotic features | 3,071 | 2,530 | 2,365 | 2,243 | 2,233 | 2,305 | 14,747 |
| Major depressive disorder, single episode, unspecified | 1,996 | 1,493 | 1,343 | 921 | 719 | 265 | 6,737 |
| Bipolar disorder, unspecified | 1,110 | 981 | 852 | 820 | 676 | 613 | 5,052 |
| Major depressive disorder, recurrent, severe with psychotic symptoms | 853 | 787 | 726 | 681 | 651 | 559 | 4,257 |
| Adjustment disorders | 635 | 604 | 822 | 830 | 601 | 378 | 3,870 |

Table 7. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents by Diagnosis¹, Age Group, and Year, 2017–2022

Note:

¹Diagnosis based on first listed mental and behavioral disorder ICD-10-CM code.

| | | | Ye | ar | | | Total |
|-----------|---------|---------|---------|---------|---------|---------|-----------|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | 10041 |
| Youth | 21,168 | 20,733 | 20,442 | 13,074 | 12,495 | 14,498 | 102,410 |
| Race | | | | | | | |
| Black | 3,361 | 3,302 | 3,254 | 1,725 | 1,495 | 2,060 | 15,197 |
| White | 17,416 | 17,030 | 16,773 | 11,069 | 10,718 | 12,011 | 85,017 |
| Other | 391 | 401 | 415 | 280 | 282 | 427 | 2,196 |
| Sex | | | | | | | |
| Female | 10,092 | 10,056 | 10,149 | 6,757 | 6,817 | 7,664 | 51,535 |
| Male | 11,076 | 10,677 | 10,293 | 6,317 | 5,678 | 6,834 | 50,875 |
| Age Group | | | | | | | |
| <10 | 4,233 | 3,980 | 3,568 | 1,999 | 1,718 | 2,129 | 17,627 |
| 10–13 | 6,141 | 6,274 | 6,370 | 3,953 | 3,792 | 4,349 | 30,879 |
| 14–17 | 10,794 | 10,479 | 10,504 | 7,122 | 6,985 | 8,020 | 53,904 |
| Adult | 217,747 | 225,504 | 234,858 | 187,750 | 170,242 | 168,873 | 1,204,974 |
| Race | | | | | | | |
| Black | 22,361 | 23,828 | 26,069 | 19,873 | 16,841 | 17,779 | 126,751 |
| White | 193,616 | 199,614 | 206,441 | 165,879 | 151,311 | 148,702 | 1,065,563 |
| Other | 1,770 | 2,062 | 2,348 | 1,998 | 2,090 | 2.392 | 12,660 |
| Sex | | | | | | | |
| Female | 145,488 | 149,959 | 157,018 | 121,760 | 110,363 | 110,371 | 794,959 |
| Male | 72,259 | 75,545 | 77,840 | 65,990 | 59,879 | 58,502 | 410,015 |
| Age Group | | | | | | | |
| 18–24 | 23,588 | 23,995 | 25,561 | 20,287 | 18,649 | 18,216 | 130,296 |
| 25–34 | 41,623 | 42,299 | 42,863 | 34,511 | 30,637 | 29,455 | 221,388 |
| 35–44 | 43,640 | 44,223 | 45,383 | 36,860 | 32,814 | 30,639 | 233,559 |
| 45–54 | 42,463 | 43,415 | 43,363 | 34,766 | 30,408 | 29,176 | 223,591 |
| 55–64 | 32,747 | 34,497 | 36,707 | 29,213 | 26,670 | 26,985 | 186,819 |
| 65–74 | 18,962 | 20,794 | 22,799 | 18,486 | 17,971 | 19,635 | 118,647 |
| 75+ | 14,724 | 16,281 | 18,182 | 13,627 | 13,093 | 14,767 | 90,674 |
| Total | 238,915 | 246,237 | 255,300 | 200,824 | 182,737 | 183,371 | 1,307,384 |

Table 8. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents: Demographics by Year, 2017–2022

Note:

| Diagnosis | | | Ye | ear | | | Total |
|---|--------|--------|--------|--------|--------|--------|---------|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | |
| Youth | | | | | | | |
| Attention-deficit/hyperactivity disorder, unspecified type | 9,196 | 8,966 | 8,289 | 4,787 | 3,686 | 4,840 | 39,764 |
| Other | 4,954 | 4,595 | 4,545 | 3,233 | 4,170 | 6,110 | 27,607 |
| Anxiety disorder, unspecified | 2,697 | 2,934 | 3,162 | 2,313 | 2,256 | 2,713 | 16,075 |
| Major depressive disorder, single episode, unspecified | 3,001 | 3,059 | 3,293 | 2,132 | 1,753 | 221 | 13,459 |
| Bipolar disorder, unspecified | 716 | 670 | 563 | 303 | 286 | 293 | 2,831 |
| Oppositional defiant disorder | 604 | 509 | 590 | 306 | 344 | 321 | 2,674 |
| Adults | | | | | | | |
| Anxiety disorder, unspecified | 71,988 | 74,577 | 77,502 | 63,889 | 58,209 | 58,806 | 404,971 |
| Major depressive disorder, single episode, unspecified | 72,301 | 74,612 | 76,411 | 56,297 | 42,351 | 4,018 | 325,990 |
| Other | 40,038 | 41,556 | 43,886 | 36,730 | 42,677 | 80,676 | 285,563 |
| Bipolar disorder, unspecified | 18,125 | 17,498 | 18,330 | 14,256 | 11,720 | 11,862 | 91,791 |
| Other specified anxiety disorders | 8,487 | 8,262 | 9,828 | 9,552 | 9,342 | 8,119 | 53,590 |
| Generalized anxiety disorder | 6,808 | 8,999 | 8,901 | 7,026 | 5,943 | 5,392 | 43,069 |

Table 9. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders among Youth and Adult Kentucky Residents by Diagnosis¹, Age Group, and Year, 2017–2022

Note:

¹Diagnosis based on first listed mental and behavioral disorder ICD-10-CM code.

| | | | Ye | ar | | | Total | | | | |
|-----------|-------|-------|-------|-------|-------|-------|--------|--|--|--|--|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | | | | | |
| Adult | | | | | | | | | | | |
| Race | | | | | | | | | | | |
| Black | 401 | 401 | 357 | 429 | 574 | 579 | 2,741 | | | | |
| White | 7,396 | 6,589 | 5,563 | 5,405 | 5,846 | 5,472 | 36,271 | | | | |
| Other | 74 | 51 | 36 | 74 | 102 | 101 | 438 | | | | |
| | | | | | | | | | | | |
| Female | 2,729 | 2,424 | 1,904 | 1,736 | 1,980 | 1,850 | 12,623 | | | | |
| Male | 5,142 | 4,617 | 4,052 | 4,172 | 4,542 | 4,302 | 26,827 | | | | |
| Age Group | | | | | | | | | | | |
| 18–24 | 565 | 444 | 280 | 273 | 293 | 263 | 2,118 | | | | |
| 25–34 | 2,152 | 1,814 | 1,445 | 1,231 | 1,408 | 1,297 | 9,347 | | | | |
| 35–44 | 2,102 | 1,922 | 1,602 | 1,675 | 1,827 | 1,710 | 10,838 | | | | |
| 45–54 | 1,671 | 1,529 | 1,295 | 1,322 | 1,423 | 1,391 | 8,631 | | | | |
| 55–64 | 1,012 | 982 | 997 | 1,018 | 1,136 | 1,035 | 6,180 | | | | |
| 65–74 | 309 | 274 | 265 | 318 | 369 | 380 | 1,915 | | | | |
| 75+ | 60 | 76 | 72 | 71 | 66 | 76 | 421 | | | | |
| Total | 7,871 | 7,041 | 5,956 | 5,908 | 6,522 | 6,152 | 39,450 | | | | |

Table 10. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders Associated with Substance Use among Kentucky Adult Residents: Demographics by Year, 2017–2022

Note:

Table 11. Numbers of Nonfatal Inpatient Hospitalizations for Mental and Behavioral Disorders Associated with Substance Use among Kentucky Adult Residents by Diagnosis¹ and Year, 2017–2022

| Diagnosis | | | Total | | | | |
|---|-------|-------|-------|-------|-------|-------|--------|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | |
| Adult | | | | | | | |
| Alcohol dependence | 3,365 | 3,193 | 3,211 | 3,460 | 3,613 | 3,631 | 20,473 |
| Other | 1,790 | 1,406 | 1,013 | 891 | 998 | 870 | 6,968 |
| Opioid addiction and dependence | 1,794 | 1,436 | 847 | 648 | 745 | 659 | 6,129 |
| Alcohol abuse | 322 | 327 | 259 | 388 | 623 | 585 | 2,504 |
| Other stimulant abuse | 286 | 392 | 363 | 335 | 337 | 230 | 1,943 |
| Other psychoactive substance use, unspecified | 314 | 287 | 263 | 186 | 206 | 177 | 1,433 |

Notes:

¹Diagnosis based on first listed mental and behavioral disorder due to substance use ICD-10-CM code.

| | Year | | | | | | | |
|-----------|-------------------------------|--------|---------|--------|--------|--------|---------|--|
| | 2017 2018 2019 2020 2021 2022 | | | | | Total | | |
| Adult | | | _ • _ • | | | | | |
| Race | | | | | | | | |
| Black | 8,711 | 8,755 | 10,066 | 9,386 | 9,681 | 9,452 | 56,051 | |
| White | 53,051 | 56,107 | 57,759 | 54,711 | 54,521 | 52,143 | 328,292 | |
| Other | 660 | 741 | 735 | 811 | 1,130 | 1,196 | 5,273 | |
| Sex | | | | | | | | |
| Female | 22,918 | 24,765 | 25,959 | 23,471 | 24,037 | 23,817 | 144,967 | |
| Male | 39,504 | 40,838 | 42,601 | 41,437 | 41,295 | 38,974 | 244,649 | |
| Age Group | | | | | | | | |
| 18–24 | 7,336 | 7,135 | 6,977 | 6,571 | 6,693 | 6,400 | 41,112 | |
| 25–34 | 16,713 | 17,464 | 17,844 | 16,737 | 16,133 | 14,664 | 99,555 | |
| 35–44 | 14,727 | 16,069 | 17,602 | 17,161 | 17,475 | 15,615 | 98,649 | |
| 45–54 | 12,418 | 12,630 | 12,813 | 12,067 | 11,894 | 11,782 | 73,604 | |
| 55–64 | 8,269 | 8,944 | 9,729 | 8,943 | 9,098 | 9,626 | 54,609 | |
| 65–74 | 2,321 | 2,592 | 2,740 | 2,622 | 3,228 | 3,618 | 17,121 | |
| 75+ | 638 | 769 | 855 | 807 | 811 | 1,086 | 4,966 | |
| Total | 62,422 | 65,603 | 68,560 | 64,908 | 65,332 | 62,791 | 389,616 | |

Table 12. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders Associated with Substance Use among Kentucky Adult Residents: Demographics by Year, 2017–2022

Note:

| Diagnosis | Year | | | | | | | |
|---|--------|--------|--------|--------|--------|--------|---------|--|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | | |
| Adult | | | | | | | | |
| Other | 22,086 | 24,135 | 25,644 | 23,912 | 26,356 | 25,290 | 147,423 | |
| Alcohol abuse | 16,489 | 16,367 | 17,127 | 15,199 | 13,647 | 13,400 | 92,229 | |
| Other psychoactive substance abuse | 10,102 | 11,637 | 10,840 | 11,451 | 10,248 | 8,125 | 62,403 | |
| Alcohol dependence | 5,901 | 6,657 | 6,797 | 6,177 | 6,017 | 5,868 | 37,417 | |
| Alcohol abuse with intoxication, unspecified | 3,705 | 3,185 | 3,816 | 4,535 | 5,298 | 7,333 | 27,872 | |
| Cannabis abuse | 4,139 | 3,622 | 4,336 | 3,634 | 3,766 | 2,775 | 22,272 | |

Table 13. Numbers of Nonfatal Emergency Department Visits for Mental and Behavioral Disorders Associated with Substance Use among Kentucky Adult Residents by Diagnosis¹ and Year, 2017–2022

Notes:

¹Diagnosis based on first listed mental and behavioral disorder due to substance use ICD-10-CM code.

| | | Total | | | | | |
|-----------|--|-------|-------|-------|-------|-------|--------|
| | Year 2017 2018 2019 2020 2021 2022 | | | | | | 10tal |
| Adult | | | | | | | |
| Race | | | | | | | |
| Black | 427 | 412 | 360 | 380 | 427 | 380 | 2,386 |
| White | 5,222 | 4,511 | 4,100 | 4,058 | 3,624 | 3,353 | 24,868 |
| Other | 60 | 58 | 54 | 58 | 64 | 78 | 372 |
| Sex | | | | | | | |
| Female | 3,083 | 2,738 | 2,427 | 2,367 | 2,163 | 1,999 | 14,777 |
| Male | 2,626 | 2,243 | 2,087 | 2,129 | 1,952 | 1,812 | 12,849 |
| Age Group | | | | | | | |
| 18–24 | 594 | 548 | 551 | 559 | 552 | 453 | 3,227 |
| 25–34 | 1,097 | 915 | 851 | 873 | 738 | 675 | 5,149 |
| 35–44 | 1,113 | 979 | 846 | 918 | 851 | 739 | 5,446 |
| 45–54 | 1,138 | 955 | 856 | 787 | 715 | 695 | 5,146 |
| 55–64 | 953 | 807 | 731 | 746 | 672 | 635 | 4,544 |
| 65–74 | 470 | 459 | 381 | 375 | 374 | 379 | 2,438 |
| 75+ | 344 | 318 | 298 | 238 | 243 | 235 | 1,676 |
| Total | 5,709 | 4,981 | 4,514 | 4,496 | 4,115 | 3,811 | 27,626 |

Table 14. Numbers of Nonfatal Drug Overdose Inpatient Hospitalizations among Kentucky Adult Residents: Demographics by Year, 2017–2022

Note:

Table 15. Numbers of Nonfatal Drug Overdose Inpatient Hospitalizations among Kentucky Adult Residents by Diagnosis¹ and Year, 2017–2022

| Diagnosis | | Year | | | | | |
|---|-------|-------|-------|-------|-------|-------|--------|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | |
| Adult | | | | | | | |
| Other | 3,249 | 2,982 | 2,675 | 2,503 | 2,177 | 2,153 | 15,739 |
| Poisoning by, adverse effect of, and underdosing of other and unspecified drugs, medicaments, and biological substances | 678 | 533 | 498 | 593 | 586 | 524 | 3,412 |
| Poisoning by, adverse effect of, and underdosing of benzodiazepines | 620 | 495 | 409 | 385 | 375 | 294 | 2,578 |
| Poisoning by, adverse effect of, and underdosing of psychostimulants | 369 | 371 | 375 | 352 | 388 | 334 | 2,189 |
| Poisoning by, adverse effect of, and underdosing of other opioids | 362 | 330 | 290 | 334 | 300 | 310 | 1,926 |
| Poisoning by, adverse effect of, and underdosing of narcotics and psychodysleptics | 431 | 270 | 267 | 329 | 289 | 196 | 1,782 |

Notes:

¹Diagnosis based on first listed drug overdose ICD-10-CM code.

| | Year | | | | | | | | |
|-----------|--------|-------|-------|--------|--------|-------|--------|--|--|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | | | |
| Adult | | | | | | | | | |
| Race | | | | | | | | | |
| Black | 857 | 885 | 839 | 1,150 | 1,115 | 1,149 | 5,995 | | |
| White | 10,670 | 8,573 | 8,170 | 9,380 | 9,226 | 7,637 | 53,656 | | |
| Other | 138 | 91 | 115 | 136 | 165 | 187 | 832 | | |
| Sex | | | | | | | | | |
| Female | 5,013 | 4,293 | 3,984 | 4,320 | 4,381 | 3,729 | 25,720 | | |
| Male | 6,652 | 5,256 | 5,140 | 6,346 | 6,125 | 5,244 | 34,763 | | |
| Age Group | | | | | | | | | |
| 18–24 | 1,674 | 1,405 | 1,221 | 1,436 | 1,425 | 1,272 | 8,433 | | |
| 25–34 | 4,008 | 3,190 | 2,998 | 3,374 | 3,031 | 2,418 | 19,019 | | |
| 35–44 | 2,703 | 2,167 | 2,208 | 2,752 | 2,793 | 2,318 | 14,941 | | |
| 45–54 | 1,666 | 1,344 | 1,266 | 1,470 | 1,542 | 1,321 | 8,609 | | |
| 55–64 | 917 | 789 | 798 | 1,006 | 1,018 | 913 | 5,441 | | |
| 65–74 | 394 | 364 | 336 | 380 | 435 | 426 | 2,335 | | |
| 75+ | 303 | 290 | 297 | 248 | 262 | 305 | 1,705 | | |
| Total | 11,665 | 9,549 | 9,124 | 10,666 | 10,506 | 8,973 | 60,483 | | |

Table 16. Numbers of Nonfatal Drug Overdose Emergency Department Visits among Kentucky Adult Residents: Demographics by Year, 2017–2022

Note:

Table 17. Numbers of Nonfatal Drug Overdose Emergency Department Visits among Kentucky Adult Residents by Diagnosis¹ and Year, 2017–2022

| Diagnosis | Year | | | | | | Total |
|---|-------|-------|-------|-------|-------|-------|--------|
| | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | |
| Adult | | | | | | | |
| Other | 3,520 | 3,469 | 3,275 | 3,311 | 3,264 | 3,094 | 19,933 |
| Poisoning by, adverse effect of, and underdosing of narcotics and psychodysleptics | 5,077 | 3,485 | 3,136 | 3,571 | 2,611 | 1,625 | 19,505 |
| Poisoning by, adverse effect of, and underdosing of other and unspecified drugs, medicaments, and biological substances | 1,314 | 1,115 | 1,166 | 1,501 | 1,851 | 1,557 | 8,504 |
| Poisoning by, adverse effect of, and underdosing of other and unspecified narcotics | 703 | 556 | 654 | 1,037 | 1,363 | 1,383 | 5,696 |
| Poisoning by, adverse effect of, and underdosing of other opioids | 517 | 458 | 514 | 803 | 970 | 1,043 | 4,305 |
| Poisoning by, adverse effect of, and underdosing of benzodiazepines | 534 | 466 | 379 | 443 | 447 | 271 | 2,540 |

Notes:

¹Diagnosis based on first listed drug overdose ICD-10-CM code.

Definitions

| | ICD-10 | /ICD-10-CM Codes |
|--|---|---|
| Indicator | Emergency Department Visits | Inpatient Hospitalizations |
| Mental and behavioral disorders | F20–F69 or F90–F99 in any field | F20–F69 and F90–F99 in the primary diagnosis field |
| Mental and behavioral disorders | F10–F16 or F18–F19 in any | F10–F16 and F18–F19 in the primary |
| due to substance use | field | diagnosis field |
| | Fatal | Nonfatal |
| Self-harm (from CSTE Injury Surveillance Toolkit) | Underlying cause of death code of X60–X84, Y87.0, or U03 | In any field: X71–X83; T36–T50 with 6th character = 2 (Note: include T36.9, T37.9, T39.9, T41.4, T42.7, T43.9, T45.9, T47.9, and T49.9 with 5th character = 2); T51–T65 with 6th character = 2 (Note: include T51.9, T52.9, T53.9, T54.9, T56.9, T57.9, T58.0, T58.1, T58.9, T59.9,T60.9, T61.0, T61.1, T61.9, T62.9, T63.9, T64.0, T64.8, and T65.9 with a 5th character = 2); T71 with 6th character = 2; and T14.91; all codes must have a 7th character of A or missing, excluding adverse effects and underdosing |
| Drug overdose (from CSTE Injury Surveillance Toolkit) | Underlying cause of death code of X40–X44, X60–X64, X85, or Y10–Y14 | In any field: T36–T50; for T36.9, T37.9, T39.9, T41.4, T42.7, T43.9, T45.9, T47.9, and T49.9, a 5th character; for all others, a 6th character; all codes must have a 7th character of A or missing, excluding adverse effects and underdosing |

Data Sources

Emergency department visits: Kentucky Outpatient Services Database, Office of Health Data and Analytics, Cabinet for Health and Family Services

Inpatient hospitalizations: Kentucky Inpatient Hospitalization Claims Files, Office of Health Data and Analytics, Cabinet for Health and Family Services

Deaths: Kentucky Death Certificate Database, Kentucky Office of Vital Statistics, Cabinet for Health and Family Services

Disclaimers

Emergency department visits and inpatient hospitalization counts represent encounters of care and could be greater than the total number of patients treated. To avoid duplicate counts, inpatient hospitalizations resulting in

death and emergency department visits resulting in inpatient hospitalization or death have been excluded from this analysis. Data are provisional and subject to change. Produced by the Kentucky Injury Prevention and Research Center as bona fide agent for the Kentucky Department for Public Health, August 2023. This report was supported by grants numbers NU17CE010064 (Centers for Disease Control and Prevention) and 3048114413. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention and Prevention or the U.S. Department of Health and Human Services.